

Parton-hadron duality in event generators

J.CHÝLA and J.RAMEŠ

*Institute of Physics, Czechoslovak Academy of Sciences
Prague, Czechoslovakia¹*

Abstract

The validity of local parton-hadron duality within the framework of HERWIG and JETSET event generators is investigated. We concentrate on e^+e^- annihilations in LEP 2 energy range as these interactions provide theoretically the cleanest condition for the discussion of this concept.

¹Postal address: Na Slovance 2, 180 40 Prague 8, Czechoslovakia

1 Introduction

The concept of parton hadron duality (PHD) and in particular its local version attempts to answer the question of the relation between the properties of experimentally observed hadrons and the assumed underlying parton dynamics. A few years ago the St.Petersburg school has gone beyond the original global version of the duality idea and has argued in favour of much closer relation between the single particle inclusive spectra of partons and hadrons [1]. They have developed powerful theoretical tools to calculate within perturbative QCD partonic spectra in great detail, taking into account various subtle effects (for comprehensive review of this topic see, for instance, [2]). In converting their results into the statements concerning hadrons they, however, crucially rely on two important assumptions. First, the independent fragmentation model is used to hadronize the partonic configurations originating from perturbative cascading. Secondly, partonic cascades are allowed to evolve down to rather small (timelike) virtualities of the order of pion mass. As neither of these assumptions is incorporated in the currently widely used generators based on either the string or cluster fragmentation, we have undertaken a detailed investigation of the PHD within two distinct event generators which successfully describe vast amount of experimental data from various collisions. For related work see also [3, 4]. Both HERWIG (we use its 5.3 version) and JETSET (version 7.2) use in their respective hadronization stages algorithms which do not allow a sensible, i.e. reasonably unambiguous interpretation of produced hadrons as being fragments of a particular single parton. In our study we have addressed two closely related questions:

- to what extent do the hadronic distributions reflect those of the perturbatively produced partons
- how much do the partonic spectra in HERWIG and JETSET differ from each other

In the local PHD picture the first question has a very simple answer, at least if the fragmentation function advocated by the authors of [1] is employed: hadronic spectra are proportional to those of the partons with the proportionality factor of the order of unity. This is not the case in either HERWIG or JETSET generator. These generators are both based on the partonic cascade as described by perturbative QCD so that the starting point is in principle the same as in [1]. The partonic cascades in HERWIG and JETSET are, however, neither identical nor fully equivalent to the analytic calculations of [1] so that differences do appear already on the parton level. Moreover, the influence of the hadronization stage turns out to be very important in its effects on the hadronic spectra. What we basically observe is that quite different configurations of partons yield much the same hadronic spectra. Indeed while HERWIG and JETSET are quite different as far as partonic spectra are concerned they yield remarkably similar results for hadrons. This indicates that the interplay between the perturbative and hadronization stages in hard scattering processes is very important, nontrivial and model dependent. Because of this ambiguity in the relation between the partonic and hadronic characteristics the concept of local PHD loses much of its intuitive appeal and predictive power.

In order to investigate these questions in relatively "clean" conditions, we have concentrated on the e^+e^- annihilations into hadrons at 200 GeV center of mass energy, i.e. on LEP 2 energy range. There the perturbative cascades, though model dependent, are already rather well developed and so the whole perturbation theory machinery seems to be well justified and under control.

Any comparison between several sets of results depends on the quantities selected for that purpose. We have chosen the following ones:

- multiplicity distribution
- single particle inclusive distributions in the standard variables z and p_t^2 where

$$z = \ln \left(\frac{1}{x_p} \right); \quad x_p = \frac{2p}{\sqrt{s}} \quad (1)$$

where p_t denotes the transversal component of particle momenta with respect to the thrust axis.

- factorial moments of particle multiplicities (intermittency measure)

The first two types of them are those discussed in [1] while the factorial moments of particle multiplicities in small regions of phase space [5, 6] are commonly considered as a clear manifestation of underlying partonic cascading [7].

2 A few remarks on generators

We briefly recall some of the important features of HERWIG and JETSET generators and in particular those of their parameters which will play an essential role in the following discussion of local PHD. In HERWIG the event simulation proceeds in four stages:

1. hard scattering, i.e. in our case $e^+e^- \rightarrow q\bar{q}$
2. evolution of parton showers on all partonic legs, here $q\bar{q}$
3. formation of colorless clusters out of partonic final state $q\bar{q}$ pairs
4. decay of these clusters into hadrons

In JETSET the first two steps are in principle similar although the details of shower evolution differ from HERWIG. The main difference between these generators concerns, however, the hadronization stage. Instead of the formation of colorless clusters JETSET spans relativistic string on the products of partonic cascade which then breaks into observable hadrons.

There are many parameters which determine the details of each of these steps, but the following ones are essential for fixing the relative importance and interplay between the parton shower and hadronization stages in HERWIG:

- *QCDLAM*: the usual QCD Λ -parameter
- *VQCUT*, *VGCUT*: parameters setting, when added to parton masses, the minimal parton virtuality in timelike cascades. The default options are such that for both quarks and gluons the minimal timelike virtuality is about 0.9 GeV.
- *CLMAX*: the decisive parameter for the hadronization stage of HERWIG. It forces the colorless clusters, produced in the perturbative cascade, with mass squared above $CLMAX^2 + (m_q + m_{\bar{q}})^2$ to split, before hadronization, into lower mass ones. The default value is 3.5 GeV. Although in typical events the action of *CLMAX* parameter is limited it plays quite an essential role in defining the relative role of partonic cascades and hadronization stages. As will be shown below it is particularly important in low parton number events.

In JETSET analogous role is played by the parameters:

- PARJ(81): analogue of *QCDLAM*
- PARJ(82): invariant mass cutoff on parton virtualities, similar in effect to *VGCUT*, *VQCUT* in HERWIG.
- there is no direct analogue of *CLMAX* parameter.

In JETSET the user can choose between the leading log parton showers and exact fixed order matrix element approaches. In HERWIG parton showers cannot be straightforwardly switched off and they always accompany the hard scattering subprocess, be it $e^+e^- \rightarrow q\bar{q}$ or $e^+e^- \rightarrow q\bar{q}g$. Nevertheless to study the relation between partons and hadrons in theoretically cleanest conditions we have concentrated in both generators on the $e^+e^- \rightarrow q\bar{q}$ subprocess and selected the parton shower option in JETSET. In our quest to understand the relation between the parton and hadron spectra we have

- compared the hadron as well as parton characteristics, specified above, for the currently "best" sets of parameters of HERWIG and JETSET
- compared parton distributions with the corresponding hadronic ones, in HERWIG as well as in JETSET
- looked separately on events characterized by different number of partons. "Tagging" on this parton number allows us to see clearly under what circumstances are the hadronic characteristics direct manifestations of the underlying partonic ones.
- varied the values of *VQCUT*, *VGCUT*, *CLMAX* and looked at the changes in the relation between partonic and hadronic spectra

3 Discussion of the results

The basic results of extensive simulations with HERWIG and JETSET at LEP2 energy are presented in Fig.1-7. We stress that neither generator has been tuned especially for our purposes and we have taken the currently "best" sets of their respective parameters.

3.1 Multiplicity distributions

Fig.1 displays the comparison between the partonic as well as hadronic multiplicity distributions in HERWIG and JETSET. Already this simplest quantity signals the basic message: while for the hadrons these models predict results which are rather close to each other (typically within 10-15 %) they differ vastly on the level of partons! Despite the large difference in the average number of perturbatively produced partons (9 in HERWIG vs. 15.5 in JETSET) the hadronic multiplicity distributions are much closer in shape as well in average values: 42.5 in HERWIG vs. 48 for JETSET. This large difference between the average parton and hadron multiplicities is in sharp contrast with the results of [1], where the number of partons is much closer to the number of hadrons.

3.2 Single particle inclusive spectra

The same message can be read off Fig.2, where the single particle spectra in z and p_t^2 , from both HERWIG and JETSET, are compared on partonic as well as hadronic levels. The pronounced difference in z distributions of partons in the region close to zero comes from the region of phase space where one of the partons carries nearly all available momentum. The most important difference between HERWIG and JETSET partonic distributions is, however, observed for large z and small p_t^2 , i.e. for soft partons which dominate the total multiplicity. We see that HERWIG z -distribution is significantly lower than that of JETSET down to $z = 2$ and practically vanishes for $z > 5$. Despite these dramatic differences on the level of partons, the corresponding hadronic distributions are quite similar even in the large z region. For p_t^2 distributions of partons the dramatic difference in the region close to $p_t^2 = 0$ is a direct manifestation of the cut on minimal virtuality of partons as set by the parameters $VQCUT, VGCUT$. The position of the maximum in HERWIG spectrum is in fact proportional to them. In JETSET there does not seem to be an analogous effect. This difference between HERWIG and JETSET is, however, again not reflected in the corresponding hadronic p_t^2 distributions, which look practically indistinguishable. The slightly higher hadronic multiplicity in JETSET is then reflected in somewhat higher values of hadronic z spectra in the region around $z = 4$.

In Fig.3 we compare the partonic and hadronic spectra obtained with both generators by plotting (as solid lines) the ratios

$$r(w) = \left(\frac{1}{N_{events}} \frac{dN}{dw_{partons}} \right) / \left(\frac{1}{N_{events}} \frac{dN}{dw_{hadrons}} \right); \quad w = z, p_t^2 \quad (2)$$

of appropriately normalized partonic over hadronic distributions, which should be approximately constant if local PHD holds. Clearly rather large deviations from constancy in

most of the phase space and for both z and p_t^2 are observed, the pattern of these violations being, except for z close to the upper limit, similar in HERWIG as in JETSET. In both generators a large part of this effect can be traced back to the fact that their partonic showers are stopped at much higher virtualities than in [1]. This is demonstrated by dotted lines in Fig.3 which correspond, for HERWIG as well as JETSET, to lower virtuality cut-off $Q_0 = 0.2$ GeV. For technical reasons this low virtuality cut-off requires, in both generators, simultaneous lowering of the QCD Λ -parameter. In our case we have taken it to be 0.04 GeV.

The most dramatic effect occurs for the ratio $r(z)$ which for $Q_0 = 1.0$ GeV was rapidly decreasing function of z in the whole phase space, while now this ratio is nearly constant in the large interval $z \in (1, 6)$. Similar effect is observed for p_t^2 spectra, in particular on low p_t^2 region. There is thus no doubt that local PHD is much better reproduced in both HERWIG and JETSET for small virtuality cut-off $Q_0 = 0.2$ GeV. However, neither HERWIG nor JETSET can accommodate such low values of Q_0 and still describe the available experimental data as accurately as with the Q_0 in the region of 1 GeV. In the regions of x_p close to 0 and 1 the deviations from constancy are very large even for $Q_0 = 0.2$ GeV but this is not surprising as these are also the regions where the analytic calculation of [1] contain subtle effects not included in the generators. An interesting difference between HERWIG and JETSET z distributions is observed in large z region, i.e. for $x_p \rightarrow 0$, which is populated by soft partons, and where the ratios behave quite oppositely. All this suggests that the validity of local PHD relies heavily on evolving the partonic showers to rather small scales comparable to the pion mass.

In order to understand the differences between HERWIG and JETSET in more detail we have furthermore subdivided all events in three classes according to the number of perturbatively produced partons.

- small number of partons: 2-6
- moderate number of partons: 7-11
- large number of partons: more than 11

In Fig.4 the comparison between partonic z and p_t^2 spectra from HERWIG and JETSET is done for each of the classes separately. We see that the differences increase with decreasing number of underlying partons. For all three classes of events the corresponding hadronic spectra (not displayed) are, however, again much closer as in the case of the full samples.

We now come to the interplay between the parton shower and cluster decay stages in HERWIG event generator. In Fig.5 we plot separately for each of the three classes of events the z and p_t^2 distributions for partons and hadrons. In the case of hadronic distributions we moreover show two sets of curves corresponding to two different values of the *CLMAX* parameter. Beside the default value $CLMAX = 3.5$ GeV we plot also the results corresponding to $CLMAX = 50$ GeV. This large value effectively means that we do not force large mass clusters to split into smaller ones before hadronization and thus come close to the original formulation of HERWIG. We conclude that

1. Small value of $CLMAX$ is more effective in events with small number of partons. For them perturbative branching itself leads on average to small number of heavy clusters. Once small $CLMAX$ is taken, all these heavy clusters are first split into smaller ones and only then allowed to hadronize. After this step the cluster mass distributions are essentially the same irrespective of the original parton number. The only trace of the originally different parton numbers then remains in the smaller cluster multiplicity.
2. The differences between the shapes as well as magnitudes of parton and hadron spectra are large in the whole phase space and depend sensitively on the number of partons.
3. The differences between the two hadronic spectra, corresponding to $CLMAX = 3.5$ GeV and $CLMAX = 50$ GeV show that for events with small parton numbers the initial part of the hadronization stage i.e. the cluster splitting is crucially important. This shows why and how the hadronization stage may significantly influence the relation between the parton and hadron spectra.

Taken together the message contained in the preceding observations is simple and clear: While on the level of hadrons different models give very similar results, these results can originate from very different partonic distributions. Moreover, the effects and importance of the hadronization stage are nontrivial and do depend on particular partonic configuration. All this demonstrates that the concept of local PHD does not hold within either HERWIG or JETSET event generators and is thus more ambiguous than as claimed in [1].

3.3 Intermittency analysis

The intermittency phenomenon, as quantified by the factorial moments in small phase space regions [5, 6] is commonly regarded as a convincing evidence for the underlying partonic cascading process with the selfsimilarity property [7]. As such it should also provide the evidence for the local PHD. To check this assumption we have carried out several simple tests using both HERWIG and JETSET event generators. The question we ask ourselves is similar as before: to what extent is the observed intermittency behaviour on the level of hadrons a direct consequence of the underlying partonic cascade?

To find the answer we have calculated the conventional factorial moments of the i -th rank in two dimensions (rapidity versus azimuthal angle)

$$F_i(y, \phi) = \frac{1}{N_{events}} \sum_{events} \frac{\sum_{k=1}^{n_{bins}} \langle n_k(n_k - 1) \cdots (n_k - i + 1) \rangle}{(\langle n \rangle / n_{bins})^i} \quad (3)$$

where $\langle n \rangle$ is the average number of particles in the full $\Delta y - \Delta \phi$ region accepted (we have taken full azimuthal coverage and $y \in (-2, 2)$) and n_{bins} denotes the number of bins in this two-dimensional space, given as 4^b with $n_{div} = 2^b$ defining the number of divisions in each of the two directions. We have constructed these moments for partons as well as hadrons as functions of b or n_{div} . Then we have compared for $i = 2, 3, 4, 5$

- the results from HERWIG and JETSET (Fig.6a)
- the results for partons with those of hadrons (Fig.6b)
- the results corresponding to events with different number of partons (Fig.7)

On the basis of the plots displayed in Figs.6,7 we draw the following conclusions, which point in the same direction as those of the preceding paragraphs:

1. Despite the large differences on the partonic level HERWIG and JETSET give very similar results for the hadronic factorial moments (3). This conclusion holds well for the full sample of events as well as for all three classes of events defined above (corresponding plots similar to Fig.6a are omitted), i.e. is independent of the number of underlying partons.
2. For HERWIG we find a remarkable agreement between the partonic and hadronic factorial moments in the region where the former show the appropriate rising behaviour. It is clear that this region can not be large as the number of partons is much smaller than that of the hadrons. In order to see the effect on the parton level clearly we have therefore chosen finer steps in the division of Δy - $\Delta\phi$ interval for partonic moments. We have, however, seen that much the same behaviour of hadronic factorial moments results even for the case where there are so little partons that their own factorial moments vanish. So again we see that the property of hadrons usually considered as a direct consequence of the presence of partonic cascade can equally well result from the effects of the hadronization stage. In other words there is some kind of duality but of different sort that originally proposed in [1], namely the duality between the partonic and hadronization stages within the whole event generation.
3. Except for F_2 the factorial moments in HERWIG as well as JETSET are practically independent of the number of partons in the event. In Fig.7 we present results from JETSET, but the same picture is obtained for HERWIG as well. For the purpose of this comparison we have somewhat changed the definition of the three classes of events (2-10,11-18,19 and more) which is more appropriate for JETSET due to its higher average parton multiplicity. This at first sight surprising observation again shows that there is no simple relation between the partonic and hadronic properties. The fact that events with small number of partons give smaller F_2 than those with many partons may be a simple reflection of lower overall hadronic multiplicity in these events. In HERWIG the approximate independence of the factorial moments has a simple explanation in the interplay between the partonic cascade and the hadronization stage. What happens is that in the case of small parton number the action of the *CLMAX* parameter creates out of the small number of heavy clusters, produced by purely perturbative cascade, much larger number of lighter ones. Instead of partonic cascade we have "cluster" cascade, which, however, produces essentially the same behaviour of hadronic factorial moments. In JETSET we can offer no such simple explanation but the effect holds as well.

4 Summary and conclusions

In this paper we have discussed the relation between partons and hadrons in two different, widely used event generators, HERWIG and JETSET. We concentrated on e^+e^- annihilations at LEP 2 energies as these conditions provide theoretically the cleanest place for investigation of such a relation. Our simulations show that this relation is complicated and generator dependent. The idea of local PHD as suggested in [1] is not realized in either of these models. To large extent, this is due to the fact that the analytical calculations of [1] rely in a crucial way on the use of independent fragmentation model coupled with the assumption that partonic showers are allowed to evolve down to virtualities of the order of the pion mass, while both HERWIG and JETSET stop their respective cascades at much higher virtualities.

On the other hand we have found evidence for the strong interplay between the effects of parton showers and those of the hadronization stage. Clear example of such an interplay is provided by factorial moments in narrow phase space region where the intermittent behaviour originates equally from perturbative parton branching as well as from colorless cluster splitting. This indicates that we can not check details of partonic evolution without the knowledge of hadronization mechanism and vice versa.

References

- [1] Ya.I.Azimov, Yu.L.Dokshitzer, V.A.Khoze, S.I.Troyan: Z. Phys. C - Particles and Fields 27 (1985) 65
- [2] Yu.L.Dokshitzer, V.A.Khoze, A.H.Mueller, S.I.Troyan: Basics of Perturbative QCD, Editions Frontieres, ed. J.Tran Thanh Van (1991)
- [3] B.Andersson, P.Dahlqvist, G.Gustafson: Z. Phys. C - Particles and Fields 44 (1989) 455
- [4] B.Andersson, P.Dahlqvist, G.Gustafson: Z. Phys. C - Particles and Fields 44 (1989) 461
- [5] A.Bialas, R.Peschanski: Nucl. Phys. B273 (1986) 703
- [6] A.Bialas, R.Peschanski: Nucl. Phys. B308 (1988) 857
- [7] W.Ochs, J.Wosiek: Phys. Lett. B214 (1988) 617

Figure captions.

Fig.1: Multiplicity distributions of partons (a) and hadrons (b) from HERWIG and JETSET.

Fig.2: Single particle distributions in z and p_t^2 of hadrons (a) and partons (b) from HERWIG and JETSET.

Fig.3: Ratia $r(z)$, $r(p_t^2)$ from JETSET (a) and HERWIG (b) for two values of the lower virtuality cut-off Q_0 .

Fig.4: Single particle distributions of hadrons from HERWIG and JETSET, as functions of z and p_t^2 , for the three classes of events defined in the text.

Fig.5: Partonic and hadronic distributions in z and p_t^2 for the three classes of events defined in the text as obtained from HERWIG. For the hadrons the spectra corresponding to two different values of $CLMAX$ parameter are displayed.

Fig.6: Comparison of factorial moments $F_i(y, \phi)$ between HERWIG and JETSET (a), and between partons and hadrons within HERWIG (b).

Fig.7: Dependence of factorial moments $F_i(y, \phi)$ on the number of partons as obtained from JETSET.